



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2016

Euthanasia or Assisted Suicide in Patients With Psychiatric Illness

Hodel, M A ; Trachsel, Manuel

DOI: <https://doi.org/10.1001/jama.2016.16283>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-127936>

Journal Article

Accepted Version

Originally published at:

Hodel, M A; Trachsel, Manuel (2016). Euthanasia or Assisted Suicide in Patients With Psychiatric Illness. JAMA : the Journal of the American Medical Association, 316(20):2153-2154.

DOI: <https://doi.org/10.1001/jama.2016.16283>

Letter to the editor

Martina A. Hodel, MSc¹; Manuel Trachsel, MD, PhD^{1,2}

¹Institute of Biomedical Ethics and Medical History, University of Zurich, Zurich, Switzerland

²Psychiatry Department, Cedars-Sinai Health System, Los Angeles, CA

Word count: 391

References: 5

Corresponding author: Manuel Trachsel, manuel.trachsel@uzh.ch

In their recent article in *JAMA Psychiatry*, Olié and Courtet¹ took the report by Kim and colleagues² on euthanasia or assisted suicide (EAS) of psychiatric patients in the Netherlands as an occasion to highlight the challenges surrounding the legalization of EAS in patients with a psychiatric illness. The present contribution comments on their propositions.

First, Olié and Courtet¹ mention the option of palliative care for terminally ill patients as a possible alternative to EAS. We believe that the achievements of contemporary palliative care indicate relevance beyond the context of terminal somatic illness, for the significant group of people suffering from severe persistent mental illness (SPMI). For this group of patients, we advocate a shift towards patient-oriented palliative care, centered on the ethical principle of patient self-determination. Given its focus on empowering people with mental illness, the recovery movement is already heading in this direction.³ However, some of those with SPMI may have a long history of failed functional recovery. Their suffering may be unbearable and their therapeutic options exhausted. In such specific cases, a reorientation of

care goals toward symptom relief as the main focus of care—possibly without modifying the course of the disease—may be a legitimate option.

Second, the authors¹ mention the possibility of EAS evaluations in the interests of suicide prevention. Although suicide prevention has a long history in psychiatry, there are potential risks in focusing mainly or exclusively on impeding suicide without taking account of the risks of overly aggressive care or even coercion. The severe side effects and interactions of polypharmacy, as well as the patient's (and doctor's) possible sense of failure, may result in an even poorer quality of life for the patient.⁴ We agree with Yager⁵ that hyper-interventionism (especially against the patient's wishes) and treatment excesses are not supportable; and that even patients with partially impaired decision-making capacity retain some capacity to make prudent decisions in respect of their long-term desires.

We are fully aware of the controversy surrounding futility and palliative care for patients suffering from SPML.^{4,5} First and foremost, however, we would observe that a palliative approach in psychiatry does not mean giving up on a patient but rather involves redefining the goals of care. This entails accepting the reality that mental illness can be fatal. With improvements in the quality of care, there might be fewer requests for EAS.

Disclosure

This work was supported by the Palliative Care Research funding program of the Swiss Academy of Medical Sciences; and the Stanley Thomas Johnson Foundation. The views expressed in this letter are those of the authors and not necessarily those of the funder. The funding body in no way influenced the authors in writing the manuscript. The authors declare to have no competing interests.

References

1. Olié E, Courtet P. The Controversial Issue of Euthanasia in Patients With Psychiatric Illness. *JAMA*. 2016;316(6):656–657.
2. Kim SY, De Vries RG, Peteet JR. Euthanasia and assisted suicide of patients with psychiatric disorders in the Netherlands 2011 to 2014. *JAMA Psychiatry*. 2016;73(4):362–368.
3. Jacobson N, Greenley D. What is recovery? A conceptual model and explication. *Psych Serv*. 2001;52:482–485.
4. Berk M, Berk L, Udina M, et al. Palliative models of care for later stages of mental disorder: maximizing recovery, maintaining hope, and building morale. *ANZJP*. 2012;46(2):92–99.
5. Yager J. The futility of arguing about medical futility in anorexia nervosa: The question is how would you handle highly specific circumstances? *AJOB*. 2015;15(7):47–50.